

## Effect of alkaline pH on bioactive molecules of epidermal mucus from *Labeo rohita* (Rahu)

[Alkali pH'nın *Labeo rohita* (Rahu) epidermal mukusundaki biyoaktif moleküllere olan etkisi]

Najiya al-Arif<sup>1</sup>, M. Sharif Mughal<sup>1</sup>, Atif Hanif<sup>2</sup>, Andleeb Batool<sup>1</sup>

<sup>1</sup>Department of Zoology, Government College University, Lahore, Pakistan.

<sup>2</sup>Quality Operation Laboratories, University of Veterinary and Animal Sciences, Lahore, Pakistan.

The original article to which this Erratum refers was published in Turk J Biochem. 36(1): 29-34. In the original published version of this article, on page 33, the reference list was containing errors starting with reference 13. Corrected reference list included below. The publisher regrets this error.

[Bu düzeltmede söz edilen makalenin kaynaklar listesinde, 33. sayfada 13. referansdan itibaren hata yapılmıştır. Düzeltilmiş kaynaklar listesi aşağıdadır. Bu hatadan dolayı özür dileriz.]

### References

- [1] Pickering AD. (1974) The distribution of mucous cells in the epidermis of the brown trout *Salmo trutta* (L) and the char *Salvelinus alpinus* (L). J Fish Biol 6:111-118.
- [2] Bly JE, Clem LW. (1992) Temperature and teleost immune mechanisms. Fish and Shellfish Immunol 2:159–171.
- [3] Fletcher TC, Grant PT. (1969) Immunoglobins in the serum and mucus of plaice, *Pleuronectes platessa*. J Biochem (Tokyo). 149:218-224.
- [4] Cameron AM, Endean R. (1973) Epidermal secretions and evolution of venom glands in fishes. Toxicon 11:401-410.
- [5] Ellis AE. (2001) Innate host defense mechanisms of fish against viruses and bacteria. Dev Comp Immunol 25(8-9):827-839.
- [6] Magnadottir B. (2006) Innate immunity of fish (overview). Fish and Shellfish Immunol 20(2):137-151.
- [7] Hellio C, Pons AM, Beaupoil C, Bourgoignon N, Gal YL. (2002) Antibacterial, antifungal and cytotoxic activities of extracts from fish epidermis and epidermal mucus. Int J Antimicrob Ag 20(3):214-219.
- [8] Rombout JHWM, Joosten EPHM. (1998) Mucosal immunity. In: Pastoret PP, Griebel P, Bazin H, Govaerts A, Editors. Handbook of vertebrate immunology. Academic Press, New York, pp. 39–40.
- [9] Patrzykat A, Gallant JW, Seo J, Pytyck J, Douglas SE. (2003) Novel Antimicrobial Peptides Derived from Flatfish Genes. Antimicrob Agents Chemother 47(8):2464-2470.
- [10] Fernandes JMO, Molle G, Kemp GD, Smith VJ. (2004) Isolation and characterisation of oncorhynchin II, a histone H1-derived antimicrobial peptide from skin secretions of rainbow trout, *Oncorhynchus mykiss*. Dev Comp Immunol 28:127–138.
- [11] Mulero I, Noga EJ, Meseguer J, Garcia-Ayala A, Mulero V. (2008) The antimicrobial peptides piscidins are stored in the granules of professional phagocytic granulocytes of fish and are delivered to the bacteria-containing phagosome upon phagocytosis. Dev Comp Immunol 32(12):1531-1538.
- [12] Silphaduang U, Colorni A, Noga E J. (2006) Evidence for widespread distribution of piscidin antimicrobial peptides in teleost fish. Diseases of Aqua Org 72(3):241-252.
- [13] Khan HA, Jhingran VG. (1975) Synopsis of Biological Data on Rohu, *Cirrhinus mrigala* (Hamilton, 1822). FAO Fisheries Synopsis No. 111, FAO, Rome. 100.14.
- [14] Schaperclaus W, Kulow H, Schreckenbach K. (1992) Fish Diseases. Oxonian Press Pvt. Ltd. New Delli, Calcutta.
- [15] Cagırgan H. (2009) Main sanitary problems and needs in Turkish fish farming. The use of veterinary drugs and vaccines in Mediterranean aquaculture. Options Méditerranéennes 86:187-195.

- [16] Bergsson G, Agerberth B, Jörnvall H, Gudmundsson GH. (2005) Isolation and identification of antimicrobial components from the epidermal mucus of Atlantic cod (*Gadus morhua*). FEBS J 272:4960–4969.
- [17] Bradford MM. (1976) A Rapid and Sensitive Method for the Quantitation of Microgram Quantities of Protein Utilizing the Principle of Protein-Dye Binding. Anal Biochem 72:248-254.
- [18] Zaika LL. (1988) Spices and herbs, their antimicrobial activity its determination. J. Food Saf. 9:97.
- [19] Ellis AE. (1990) Lysozyme assays. In: Techniques In Fish Immunology (Stolen JS, Fletcher T C, Anderson DP, Roberson BS, Van Muiswinkel WB. eds.). Fair Haven, NJ: SOS Publications:101-103.
- [20] Laemmli UK. (1970) Cleavage of structural proteins during the assembly of bacteriophage T4. Nature 227:680-685.
- [21] Mozumder MH. (2005) Antibacterial activity in fish mucus from farmed fish. M.Sc Thesis. Department of Marine Biotechnology Norwegian College of Fishery Science University of Tromsø, Norway; p.20-21.
- [22] Cho S, Lee B, An H, Eun J. (2005) Kenojeinin I, antimicrobial peptide isolated from the skin of the fermented skate, *Raja kenojei*. Peptides 26:581-587.
- [23] Fernandes JMO, Kemp GD, Molle MG, Smith VJ. (2002) Antimicrobial properties of histone H2A from skin secretions of rainbow trout, *Oncorhynchus mykiss*. J Biochem 368:611–620.
- [24] Kitani Y, Tsukamoto C, Zhang G, Nagai H, Ishida M, Ishizaki S, Shimakura K, Shiomi K, Nagashima Y. (2007) Identification of an antibacterial protein as L-amino acid oxidase in the skin mucus of rockfish *Sebastes schlegeli*. FEBS J 274(1):125-136.
- [25] Subramanian S, Ross NW, MacKinnon SL. (2008) Comparison of antimicrobial activity in the epidermal mucus extracts of fish. Comp Biochem Physiol B Biochem Mol Biol 150(1):85-92.
- [26] Lemaitre C, Orange N, Saglio, P, Saint, N, Gagnon J, Molle G. (1996) Characterization and ion channel activities of novel antimicrobial proteins from the skin mucosa of carp (*Cyprinus carpio*). J Biochem 240:143-149.
- [27] Smith VJ, Fernandes JMO, Jones S J, Kemp GD, Tatner MF. (2000) Antimicrobial proteins in rainbow trout, *Oncorhynchus mykiss*. Fish and Shellfish Immun 10:243-260.