


Amentoflavone protects against hydroxyl radical-induced DNA damage via antioxidant mechanism

[Amentoflavon antioksidan mekanizma ile hidroksil radikali ile indüklenmiş DNA hasarına karşı koruyucu etkiye sahiptir]*

Xican Li^{1,**}, Li Wang^{1,**}, Weijuan Han¹, Wenqiong Mai¹, Lu Han¹, Dongfeng Chen²

¹School of Chinese Herbal Medicine, ²School of Basic Medical Science, Guangzhou University of Chinese Medicine, Guangzhou, 510006, China

Supplementary File 1. The structures of the major oxidative DNA lesions



References

- [S1] Halliwell B, Gutteridge JM, Aruoma OI. The deoxyribose method: a simple "test-tube" assay for determination of rate constants for reactions of hydroxyl radicals. *Anal Biochem* 1987; 165: 215-219.
- [S2] Valavanidis A, Vlachogianni T, Fiotaks C. 8-Hydroxy-2'-deoxyguanosine (8-OHdG): A critical biomarker of oxidative stress and carcinogenesis. *J of Environ Sci Health C Environ Carcinog Ecotoxicol Rev* 2009; 27: 120-139.
- [S3] Suzuki S, Tannka T, Poyurovsky MV, Nagano H, Mayama T, Ohkubo S, Lolshin M, Hosokawa H, Nakayama T, Suzuki Y, Sugano S, Sato E, Nagao T, Yokote K, Tatsuno I, Prives C. Phosphate-activated glutaminase (GLS2), a p53-inducible regulator of glutamine metabolism and reactive oxygen species. *Proc Natl Acad Sci USA* 2010; 107: 7461-7466.
- [S4] Shigenaga MK, Ames BN. Urinary 8-hydroxy-2'-deoxyguanosine as a biological marker of in vivo oxidative DNA damage. *Proc Natl Acad Sci USA* 1989; 86: 9697-9701.
- [S5] Cooke MS, Evans MD. 8-Oxo-deoxyguanosine: reduce, reuse, recycle? *Proc Natl Acad Sci USA* 2007; 104: 13535-13536.
- [S6] Wagner JR, Hu CC. Endogenous oxidative damage of deoxycytidine in DNA. *Proc Natl Acad Sci USA* 1992; 89: 3380-3384.
- [S7] Kreuzer DA, Essigmann JM. Oxidized, deaminated cytosines are a source of C→T transitions in vivo. *Proc Natl Acad Sci USA* 1992; 95: 3578-3582.
- [S8] Malins DC, Polissart NL, Gunselman SJ. Progression of human breast cancers to the metastatic state is linked to hydroxyl radical-induced DNA damage. *Proc Natl Acad Sci USA* 1996; 93: 2557-2563.
- [S9] Cheng KC, Cahill DS, Kasai H, Nishimura S, Loeb LA. 8-Hydroxyguanine, an abundant form of oxidative DNA damage, causes G →T and A→C substitutions, *J Biol Chem* 1992; 267: 166-172.
- [S10] Fang YZ, Zheng RL. *Reactive oxygen species in vivo in Theory and application of free radical biology*. 2nd, Science Press, Beijing China, 2002; pp. 142.
- [S11] Denq RY, Fridovich I. Formation of endonuclease III-sensitive sites as a consequence of oxygen radical attack on DNA. *Free Radical Biol Med* 1989; 6: 123-129.
- [S12] Fraga CG, Shigenaga MK, Park JW, Degan P, Ames BN. Oxidative damage to DNA during aging: 8-hydroxy-2'-deoxyguanosine in rat organ DNA and urine. *Proc Natl Acad Sci USA* 1990; 87: 4533-4537.
- [S13] Jaruga P, Rozalski R, Jawien A, Migdalski A, Olinski R, Dizdaroğlu M. DNA Damage products (5'R)- and (5'S)-8,5'-cyclo-2'-deoxyadenosines as potential biomarkers in human urine for atherosclerosis. *Biochemistry* 2012; 51: 1822-1824.
- [S14] Kuraoka I, Robins P, Masutani C, Hanaoka F, Gasparutto D, Cadet J, Wood RD, Lindahl T. Oxygen free radical damage to DNA. Translesion synthesis by human DNA polymerase η and resistance to exonuclease action at cyclopurine deoxynucleoside residues. *J Biol Chem* 2001; 276: 49283-8.